arXiv:1602.05884v1 [math.GT] 18 Feb 2016

AN EXPLICIT RELATION BETWEEN KNOT GROUPS IN LENS SPACES AND THOSE IN S^3

YUTA NOZAKI

Abstract. We consider a cyclic covering map $(\Sigma, K) \to (\Sigma', K')$ between pairs of a 3-manifold and a knot, and describe the fundamental group $\pi_1(\Sigma \setminus K)$ in terms of $\pi_1(\Sigma' \setminus K')$. As a consequence, we give an alternative proof for the fact that a certain knot in S^3 cannot be represented as the preimage of any knot in a lens space. In our proofs, the subgroup of a group G generated by the commutators and the pth power of each element of G plays a key role.

Contents

1. Introduction 1
2. Lemmas in group theory and knot theory 3
3. Main theorem and corollaries 5
Appendix A. Symmetric groups and related contents 8
Appendix B. Taking C^p repeatedly 9
References 10

1. Introduction

Let K be a torus knot and $p \geq 2$ be an integer. Hartley [7] created a decision argument to know whether K can be represented as the preimage of a knot in the lens space $L(p, q)$ for some q. For example, the trefoil $T_{3,2}$ does not appear as the preimage of any knot in $L(2, 1) \cong \mathbb{R}P^3$. The Alexander polynomial of a torus knot was used in his proof.

Let K be a knot such that the outer automorphism group $\mathrm{Out}(\pi_1(S^3 \setminus K))$ is trivial. For instance, 9_{32}, 9_{33} and 24 more prime knots with 10 crossings (and their mirror images) satisfy this condition (see Kawauchi [11, Appendix F.2] or Kodama-Sakuma [12, Table 3.1]). Then it follows from Borel's theorem (see Conner-Raymond [6, Theorem 3.2]) that K cannot be represented as the preimage of any knot in any lens space (see [11, Theorems 10.6.2 and Theorem 10.6.6 (1)]).

The purpose of this paper is to deduce the above facts from a single result. Let G be a group and $p \geq 1$ be an integer. (p is not necessarily

2010 *Mathematics Subject Classification.* 57M25, 57M27.
Key words and phrases. Knot group, lens space, covering space, pth power.

prime.) We introduce the subgroup $\mathrm{C}^p(G)$ generated by the commutators and the pth power of each element of G. The case of $p = 2$ was studied, for example, in Sun [19] and Haugh-MacHale [9], from a purely algebraic point of view. Moreover, the case where p is prime can be found in Stallings [18] and Cochran-Harvey [4] (see Remark 2.2 and Appendix B). We apply C^p to the fundamental group of the complement of a knot in a 3-manifold. Let $\pi\colon \Sigma \to \Sigma'$ be a p-fold cyclic covering, where Σ is an integral homology 3-sphere. The main result of this paper is the following.

Theorem 1 (Theorem 3.1)**.** Let K' be a knot in Σ' with the connected preimage $K := \pi^{-1}(K')$. Then the image of $\pi_*\colon \pi_1(\Sigma \setminus K) \rightarrowtail \pi_1(\Sigma' \setminus K')$ coincides with $\mathrm{C}^p(\pi_1(\Sigma' \setminus K'))$.

Considering the case of $\Sigma = S^3$, $\Sigma' = L(p, q)$, we obtain the following corollaries as mentioned before.

Corollary 1 (Corollary 3.4)**.** Let $m, n, p \in \mathbb{Z}_{\geq 2}$ with $\gcd(m, n) = 1$. There exists an integer q and a knot K' in $L(p, q)$ such that $\pi^{-1}(K')$ is isotopic to the torus knot $T_{m,n}$ or its mirror image if and only if $\gcd(mn, p) = 1$.

Furthermore, Chbili [3] determined the above q, which also follows from Corollary 1 and some other results (see Proposition 3.6 for details).

Corollary 2 (Corollary 3.3)**.** A knot K in S^3 with $\mathrm{Out}(\pi_1(S^3 \setminus K)) = 1$ cannot be represented as the preimage of any knot in any lens space.

Another proof of a special case of Corollary 2 is given in Appendix A.1. We expect that the use of Theorem 1 in the appendix can be extended to some other knots.

When a knot K is represented as a preimage, there is still the question of whether K' is uniquely determined. Sakuma [16], Boileau and Flapan [1] independently proved that if the preimages of oriented knots K'_0, K'_1 are the same prime knot, then there exists a diffeomorphism $(L(p, q), K'_0) \to (L(p, q), K'_1)$ preserving the orientations both of $L(p, q)$ and K_i. Note that K'_0 and K'_1 are not necessarily ambient isotopic to each other. (They also showed that the same is true for a composite knot under a condition regarding "slopes".) Furthermore, Manfredi [13] recently constructed two knots in a lens space such that they are not ambient isotopic to each other but their preimages are the unknot.

Our work is closely related to freely periodic actions on (S^3, K). Indeed, such an action of period p induces a cyclic covering $(S^3, K) \to (L(p, q), K')$ for some q. Hartley [7] gave a list of possible free periods of prime knots up to 10 crossings.

Acknowledgments. The author would like to express his gratitude to Takuya Sakasai for his various suggestions. Also, he would like to thank Makoto Sakuma for pointing out some mistakes in his first draft and answering his questions about the symmetry of knots. Finally, this work was supported by the Program for Leading Graduate Schools, MEXT, Japan.

2. Lemmas in group theory and knot theory

First of all, we set up notation and prepare basic lemmas used in Section 3.

2.1. **Notation.** Let X be a topological space and G be a group. We set $H_*(X) := H_*(X;\mathbb{Z})$ and $H_*(G) := H_*(G;\mathbb{Z})$ with the trivial action $G \curvearrowright \mathbb{Z}$. For a knot K in S^3, let $G(K)$ denote the fundamental group of the complement of K, that is, $G(K) = \pi_1(S^3 \setminus K)$. (We omit a base point if we need not to care it.) We denote by $\mathbb{Z}_n$ the cyclic group of order n.

2.2. **Definition of C^p and related lemmas.** We first introduce an important subgroup and give some examples.

Definition 2.1. For a group G and $p \in \mathbb{Z}_{\geq 1}$, let $\mathrm{C}^p(G)$ denote the subgroup of G generated by the set $\{g^p \mid g \in G\} \cup \{[g,h] \mid g,h \in G\}$, where $[g,h] := ghg^{-1}h^{-1}$. A group G is called a *C^p-group* if there is a group G' such that $G \cong \mathrm{C}^p(G')$.

Note that $\mathrm{C}^1(G) = G$ and $\mathrm{C}^2(G) = \langle \{g^2 \mid g \in G\} \rangle$.

Remark 2.2. $\mathrm{C}^2(G)$ is denoted by G^2 in [19] and by $S(G)$ in [9]. For a prime p, $\mathrm{C}^p(G)$ coincides with the 1st term of the p-lower central series in [18] and with the 1st term of the derived p-series in [4].

Example 2.3. Let us determine the subgroup $\mathrm{C}^p\mathfrak{S}_n$ of the nth symmetric group $\mathfrak{S}_n$. In the case where p is odd, $\mathrm{C}^p\mathfrak{S}_n$ contains all transpositions, and thus $\mathrm{C}^p\mathfrak{S}_n = \mathfrak{S}_n$. In the case where p is even, we conclude $\mathrm{C}^p\mathfrak{S}_n = \mathfrak{A}_n$ since $\mathfrak{A}_n = [\mathfrak{S}_n, \mathfrak{S}_n] \leq \mathrm{C}^p\mathfrak{S}_n \leq \mathfrak{A}_n$, where $\mathfrak{A}_n$ is the nth alternating group.

We next state two basic lemmas without proofs.

Lemma 2.4. *For any group G, the subgroup $\mathrm{C}^p(G)$ is characteristic, and the quotient group $G/\mathrm{C}^p(G)$ is abelian. Moreover, if G is finitely generated, then $G/\mathrm{C}^p(G)$ is isomorphic to a finite direct sum of finite cyclic groups whose orders are divisors of p.*

Some simple examples of $G/\mathrm{C}^n(G)$ are listed in Table 2.2.

G	$G/\mathrm{C}^p(G)$
$\mathbb{Z}_n$	$\mathbb{Z}_{\gcd(n,p)}$
$(\mathbb{Q}_{>0}, \times)$	$(\mathbb{Z}_p)^{\oplus\{\text{prime numbers}\}}$
$(\mathbb{R}_{>0}, \times)$	0

Table 2.1. Examples of $G/\mathrm{C}^p(G)$. Note that $\gcd(0,p) = p$.

Lemma 2.5. *For groups G and H, the following hold.*

1. $\mathrm{C}^p(G \times H) = \mathrm{C}^p(G) \times \mathrm{C}^p(H)$.
2. *A homomorphism $f\colon G \to H$ induces the homomorphism $\mathrm{C}^p(G) \to \mathrm{C}^p(H)$ denoted by $\mathrm{C}^p(f)$. Moreover, if f is surjective (resp. injective), then $\mathrm{C}^p(f)$ is also surjective (resp. injective).*

The following lemma is a refinement of the well-known fact [14, 13.5.8] for a complete group H.

Lemma 2.6. *Let G, H be groups such that $H \lhd G$ and $\mathrm{Ad}_g|_H \in \mathrm{Inn}(H)$ for any $g \in G$. Then the sequence of groups*

$$1 \to Z(H) \xrightarrow{\phi} H \times C_G(H) \xrightarrow{\psi} G \to 1$$

is exact, where $C_G(H)$ is the centralizer of H in G, and $\phi(h) := (h, h^{-1})$, $\psi(h, g) := hg$.

The next lemma follows from the above lemmas, which is a generalization of [9, Theorem 1].

Lemma 2.7. *Let G, H be as in Lemma 2.6 and suppose $\mathrm{C}^p(G) = H$, $Z(H) = 1$. Then $\mathrm{C}^p(H) = H$.*

Example 2.8. Let us determine whether $\mathfrak{S}_n$ is a C^p-group or not. By Example 2.3, if p is odd or $n = 1, 2$, then $\mathfrak{S}_n$ is a C^p-group. In the other cases except $n = 6$, $\mathfrak{S}_n$ is not a C^p-group by Lemma 2.7 and the fact that $\mathfrak{S}_n$ is complete for $n \neq 2, 6$. Furthermore, $\mathfrak{S}_6$ is not a C^p-group (see Appendix A.2).

Lemma 2.9. *For a group G whose abelianization is isomorphic to $\mathbb{Z}^{\oplus r}$, the quotient group $G/\mathrm{C}^p(G)$ is isomorphic to $(\mathbb{Z}_p)^{\oplus r}$.*

Proof. Since the codomain of the projection $G \twoheadrightarrow G/\mathrm{C}^pG$ is abelian, it factors through the abelianization $\mathbb{Z}^{\oplus r}$ of G. Hence, $G/\mathrm{C}^pG \cong \bigoplus_{i=1}^r \mathbb{Z}_{p_i}$ for some $p_i \mid p$. The abelianization $G \twoheadrightarrow \mathbb{Z}^{\oplus r}$ induces a surjection $G/\mathrm{C}^pG \twoheadrightarrow \mathbb{Z}^{\oplus r}/\mathrm{C}^p(\mathbb{Z}^{\oplus r}) = (\mathbb{Z}_p)^{\oplus r}$. It follows that $p_i = p$ for $i = 1, \ldots, r$. □

Remark 2.10. If p is prime, then Lemma 2.9 directly follows from [18, Theorem 3.4] or [4, Lemma 2.3].

2.3. **Links in a rational homology 3-sphere.** The following lemma is a special case of [8, Theorem 2.1].

Lemma 2.11. *Let L' be an r-component link in a rational homology 3-sphere Σ' such that $H_1(L') \xrightarrow{\mathrm{incl}_*} H_1(\Sigma')$ is surjective. Then the following holds:*

$$H_*(\Sigma' \setminus L') \cong \begin{cases} \mathbb{Z} & \text{if } * = 0, \\ \mathbb{Z}^{\oplus r} & \text{if } * = 1, \\ \mathbb{Z}^{\oplus (r-1)} & \text{if } * = 2, \\ 0 & \text{otherwise.} \end{cases}$$

Remark 2.12. If $H_1(\Sigma') \cong \mathbb{Z}_p$ and $r = 1$, then a meridian of a knot L' corresponds to $\pm p \in \mathbb{Z} \cong H_1(\Sigma' \setminus L')$.

Remark 2.13. Let Σ', L' be as in Lemma 2.11. Suppose that the universal cover of Σ' is S^3, and the link $\pi^{-1}(L')$ in S^3 is non-split. Then $H_*(\pi_1(\Sigma' \setminus L'))$ is isomorphic to $H_*(\Sigma' \setminus L')$. Indeed, since the total space $S^3 \setminus \pi^{-1}(L')$ is an Eilenberg-MacLane space, so the base space $\Sigma' \setminus L'$ is.

3. Main theorem and corollaries

Recall that we are interested in knowing whether a knot K in S^3 is represented as the preimage of a knot K' in a lens space.

3.1. **Application of C^p to covering spaces.** Let $\pi\colon \Sigma \to \Sigma'$ be a p-fold cyclic covering, where Σ is an integral homology 3-sphere. One can check that $H_*(\Sigma') \cong H_*(L(p,q))$.

Theorem 3.1. *Let K' be a knot in Σ' with the connected preimage $K := \pi^{-1}(K')$. Then the image of $\pi_*\colon \pi_1(\Sigma \setminus K) \rightarrowtail \pi_1(\Sigma' \setminus K')$ coincides with $\mathrm{C}^p(\pi_1(\Sigma' \setminus K'))$.*

Proof. Set $G := \pi_1(\Sigma \setminus K)$ and $G' := \pi_1(\Sigma' \setminus K')$. Since there is a surjection $(G')^{\mathrm{ab}} \twoheadrightarrow G'/\mathrm{C}^pG'$, by Lemma 2.11, $G'/\mathrm{C}^pG' \cong \mathbb{Z}_{p'}$ for some $p' \mid p$.

On the other hand, the covering map π induces an exact sequence

$$1 \to G \xrightarrow{\pi_*} G' \to \mathbb{Z}_p \to 1.$$

Using $\mathrm{C}^pG \le \pi_*(G) \le G'$, we have

$$p' = [G' : \mathrm{C}^pG'] = [G' : \pi_*(G)][\pi_*(G) : \mathrm{C}^pG'] \ge p.$$

It follows that $p' = p$, and thus $\pi_*(G) = \mathrm{C}^pG'$. □

Remark 3.2. Let r be the number of connected components of $\pi^{-1}(K')$ and fix the isomorphism $H_1(\Sigma') \cong \mathbb{Z}_p$ so that $[K']$ corresponds to $\overline{k'}$, where $k' := p/\mathrm{ord}([K']) \in \mathbb{Z}$. Then we have $r = k'$. In particular, $\pi^{-1}(K')$ is connected if and only if $[K']$ generates $H_1(\Sigma)$. Indeed, we first obtain the canonical isomorphism $H_1(\Sigma') \cong \pi_1(\Sigma')/\pi_*\pi_1(\Sigma)$ since the projection $\pi_1(\Sigma') \twoheadrightarrow \pi_1(\Sigma')/\pi_*\pi_1(\Sigma) \cong \mathbb{Z}_p$ factors through $\pi_1(\Sigma')^{\mathrm{ab}}$. Choose a connected component K of $\pi^{-1}(K')$ and a base point $* \in K$, and set $*' := \pi(*)$. (The other connected components are $\tau^r(K), \tau^{2r}(K), \dots, \tau^{p-r}(K)$, where τ is a generator of the deck transformation group.) It follows from $\pi(K) = K'^{p/r}$ as based loops that $\frac{p}{r} \times \overline{k'} = 0$, and thus $r \mid k'$. On the other hand, the equality $\frac{p}{k'} \times [K'] = 0$ implies that $[K']^{p/k'} \in \pi_1(\Sigma', *')$ belongs to the image of π_*. Hence, $K'^{p/k'}$ can be lifted to Σ. Then we conclude $\frac{p}{r} \mid \frac{p}{k'}$, and thus $k' \mid r$.

The case of $\Sigma = S^3$ is discussed in [13, Proposition 2.2].

As mentioned in Section 1, the following corollary is already known. However, using Theorem 3.1, we give an alternative proof of it.

Corollary 3.3. *A knot K in S^3 with $\mathrm{Out}(G(K)) = 1$ is not represented as the preimage of any knot in any lens space.*

Proof. Since $\mathrm{Out}(G(T_{m,n})) \cong \mathbb{Z}_2$ is proven in [17], K is not a torus knot, and thus $Z(G(K)) = 1$ by [2]. Hence, $G(K)$ is complete.

Assume that there exists a knot K' in $L(p,q)$ whose preimage is isotopic to K, and set $G' := \pi_1(L(p,q) \setminus K')$. Then we have $G(K) = \mathrm{C}^pG'$ by Theorem 3.1. Since $G(K)$ is complete, by Lemme 2.7, we conclude $\mathrm{C}^pG' = G'$. However, this contradicts Lemmas 2.11 and 2.9. □

3.2. **Application of Theorem 3.1 to torus knots.** The aim of this subsection is to prove the following corollary.

Corollary 3.4 ([7, Theorem 3.1])**.** *Let $m, n, p \in \mathbb{Z}_{\geq 2}$ with $\gcd(m,n) = 1$. There exists an integer q and a knot K' in $L(p,q)$ such that $\pi^{-1}(K')$ is isotopic to the torus knot $T_{m,n}$ or its mirror image if and only if $\gcd(mn,p) = 1$.*

We have to prepare the next lemma in homology of groups.

Lemma 3.5. *Let m, n, p be as in Corollary 3.4 and G be a group admitting the following exact sequence*

$$1 \to \mathbb{Z}_m * \mathbb{Z}_n \hookrightarrow G \to \mathbb{Z}_p \to 1. \tag{3.1}$$

If $\gcd(mn,p) = 1$, then $H_*(G) \cong H_*(\mathbb{Z}_{mnp})$. Moreover, the converse is true under the additional condition $\mathrm{C}^p(G) = \mathbb{Z}_m * \mathbb{Z}_n$.

Proof. Let $g \in G$ and we want to show that $\mathrm{Ad}_g|_{\mathbb{Z}_m * \mathbb{Z}_n} \in \mathrm{Inn}(\mathbb{Z}_m * \mathbb{Z}_n)$. It is known that $\mathrm{Aut}(\mathbb{Z}_m * \mathbb{Z}_n)$ is generated by inner automorphisms and $\phi_{i,j}$ for $i \in (\mathbb{Z}_m)^\times$, $j \in (\mathbb{Z}_n)^\times$ where $\phi_{i,j}$ is defined by $\phi_{i,j}(a) = a^i$, $\phi_{i,j}(b) = b^j$. Therefore, $\mathrm{Ad}_g|_{\mathbb{Z}_m * \mathbb{Z}_n} = \mathrm{Ad}_w \circ \phi_{i,j}$ for some $1 \leq i < m$, $1 \leq j < n$ and $w \in \mathbb{Z}_m * \mathbb{Z}_n$, and then the induced automorphism $(\mathrm{Ad}_g|_{\mathbb{Z}_m * \mathbb{Z}_n})^{\mathrm{ab}}$ sends n, m to in, jm respectively. On the other hand, $(\mathrm{Ad}_g)^{\mathrm{ab}} = \mathrm{id}_{\mathbb{Z}_{mnp}}$.

Let us consider the five-term exact sequence

$$0 = H_2(\mathbb{Z}_p) \to (\mathbb{Z}_{mn})_{\mathbb{Z}_p} \to \mathbb{Z}_{mnp} \to \mathbb{Z}_p \to 0$$

for the short exact sequence (3.1). Comparing the order of each group, we conclude that $(\mathbb{Z}_{mn})_{\mathbb{Z}_p} = \mathbb{Z}_{mn}$. It follows that $in = n$ and $jm = m$ in $\mathbb{Z}_{mn}$, and thus $i = j = 1$. Hence, the action $G \curvearrowright H_*(\mathbb{Z}_m * \mathbb{Z}_n)$ is trivial, and so is $\mathbb{Z}_p \curvearrowright H_*(\mathbb{Z}_m * \mathbb{Z}_n)$.

The condition $\gcd(mn,p) = 1$ implies that the E^2-term of the Lyndon-Hochschild-Serre spectral sequence for (3.1) is as follows:

$$E^2_{s,t} = H_s(\mathbb{Z}_p; H_t(\mathbb{Z}_m * \mathbb{Z}_n)) = \begin{cases} \mathbb{Z} & \text{if } s = t = 0, \\ \mathbb{Z}_{mn} & \text{if } s = 0,\ t \geq 1 \text{ is odd}, \\ \mathbb{Z}_p & \text{if } s \geq 1 \text{ is odd},\ t = 0, \\ 0 & \text{otherwise}, \end{cases}$$

and then all differentials d^r are zero for $r \geq 2$. Using $\gcd(mn,p) = 1$ again, we have $H_*(G) \cong H_*(\mathbb{Z}_{mnp})$.

We next prove the latter of the above statement. Suppose $\mathrm{C}^p(G) = \mathbb{Z}_m * \mathbb{Z}_n$. Since we know that $\mathrm{Ad}_g|_{\mathbb{Z}_m * \mathbb{Z}_n} \in \mathrm{Inn}(\mathbb{Z}_m * \mathbb{Z}_n)$ and $Z(\mathbb{Z}_m * \mathbb{Z}_n) = 1$, by Lemma 2.7, we have $\mathrm{C}^p(\mathbb{Z}_m * \mathbb{Z}_n) = \mathbb{Z}_m * \mathbb{Z}_n$. Finally, the abelianization induces a surjection $\mathbb{Z}_m * \mathbb{Z}_n / \mathrm{C}^p(\mathbb{Z}_m * \mathbb{Z}_n) \twoheadrightarrow \mathbb{Z}_{mn} / \mathrm{C}^p \mathbb{Z}_{mn} = \mathbb{Z}_{\gcd(mn,p)}$, and thus $\gcd(mn,p) = 1$. □

Note that the following proof does not assert that if $\gcd(mn,p) \neq 1$, then $G(T_{m,n})$ is not a C^p-group.

Proof of Corollary 3.4. If $\gcd(mn, p) = 1$, then a construction of a desired knot K' is given in [7, Theorem 3.1].

Suppose there exists K' as in the statement, and set $\pi_1' := \pi_1(L(p,q)\setminus K')$. The covering map $\pi\colon S^3 \setminus T_{m,n} \to L(p,q) \setminus K'$ induces the exact sequence

$$1 \to G(T_{m,n}) = \langle a, b \mid a^m = b^n \rangle \xrightarrow{\pi_*} \pi_1' \to \mathbb{Z}_p \to 1.$$

Since the center $Z(G(T_{m,n})) = \langle a^m \rangle = \mathbb{Z}$ is characteristic in $G(T_{m,n})$, the subgroup $N := \pi_*(\langle a^m \rangle)$ of π_1' is normal. We deduce the exact sequence

$$1 \to \langle a, b \mid a^m = 1 = b^n \rangle \xrightarrow{\pi_*} \pi_1'/N \to \mathbb{Z}_p \to 1$$

from the 3rd isomorphism theorem. The group $G := \pi_1'/N$ admits (3.1) and, by Theorem 3.1, it satisfies

$$\mathrm{C}^p(G) = \mathrm{C}^p(\pi_1')/N = G(T_{m,n})/\langle a^m \rangle = \mathbb{Z}_m * \mathbb{Z}_n.$$

Hence, by Lemma 3.5, it suffices to prove $H_*(G) \cong H_*(\mathbb{Z}_{mnp})$.

By Lemma 2.11 and Remark 2.13, the five-term exact sequence for

$$1 \to N \to \pi_1' \to G \to 1 \tag{3.2}$$

is as follows:

$$0 \to H_2(G) \to \mathbb{Z}_G \xrightarrow{\phi} \mathbb{Z} \to H_1(G) \to 0.$$

Since the generator a^m of N is decomposed as $\left(\prod_{i=1}^n g_i \mu g_i^{-1}\right)^m$ in π_1' for a meridian μ of K' and some $g_i \in \pi_1'$, we conclude $\phi(\overline{1}) = mnp$ by Remark 2.12. Therefore, $\mathbb{Z}_G = \mathbb{Z}$ (namely, $G \curvearrowright H_1(N)$ is trivial), $H_2(G) = 0$ and $H_1(G) = \mathbb{Z}_{mnp}$. The E^2-term of the Lyndon-Hochschild-Serre spectral sequence for (3.2) is as follows:

$$E^2_{s,t} = H_s(G; H_t(N)) = \begin{cases} H_s(G; H_0(N)) & \text{if } t = 0, \\ H_s(G) & \text{if } t = 1, \\ 0 & \text{otherwise.} \end{cases}$$

Moreover, the differential $d^2_{s,0}$ is an isomorphism for $s \geq 3$, and $H_s(G; H_0(N)) = H_0(N)_G$ is isomorphic to $\mathbb{Z}$. Therefore, the action $G \curvearrowright H_0(N)$ is trivial, and thus $H_s(G; H_0(N)) = H_s(G)$. We inductively conclude that $H_1(G) \cong H_3(G) \cong \cdots$ and $H_2(G) \cong H_4(G) \cong \cdots$. □

We finally discuss an integer q in Corollary 3.4.

Proposition 3.6 ([3, Remark 3.7])**.** *Let $|m|, |n|, p \in \mathbb{Z}_{\geq 2}$ with $\gcd(m,n) = 1$. Suppose that the torus knot $T_{m,n}$ is isotopic to the preimage of a knot in $L(p,q)$ for some q. Then $q \equiv m(1-pp^*)/n \mod p$, where p^* is an integer satisfying $pp^* \equiv 1 \mod n$.*

Proof. We first remark that $\gcd(mn, p) = 1$ by Corollary 3.4. Hence, p^* exists and $\gcd(p, m(1-pp^*)/n) = \gcd(p, 1-pp^*) = 1$ holds.

Since $T_{m,n}$ is a prime knot, by [16, Theorem 5] or [1, Theorem 1], q is uniquely determined up to taking its inverse in $\mathbb{Z}_p$. (Recall that $L(p,q_0) \cong L(p,q_1)$ as oriented manifolds if and only if $q_0 \equiv q_1^{\pm 1} \mod p$.) Therefore, it

suffices to see that there exists a knot in $L(p,q)$ whose preimage is $T_{m,n}$ for $q = m(1-pp^*)/n$. Let $q := m(1-pp^*)/n$. We now conclude $p \mid m - nq$. Hence, using a suitable n-braid (see [13, Proposition 3.3]), we construct a desired knot in $L(p,q)$. □

Appendix A. Symmetric groups and related contents

A.1. **Another proof of Corollary 3.4 for the trefoil.** The purpose of this subsection is to introduce another use of Theorem 3.1. We first remark that a quotient group of a C^p-group G is not a C^p-group in general. However, if $H \lhd G$ is characteristic, then G/H is also a C^p-group, that is, the following lemma holds.

Lemma A.1 (see [19, Theorem 1])**.** *Let G be a C^p-group and $G \twoheadrightarrow G'$ be a surjective homomorphism whose kernel is a characteristic subgroup of G. Then G' is also a C^p-group.*

Theorem A.2. *The trefoil $T_{3,2}$ cannot be represented as the preimage of a knot in $L(p,q)$ for any $p \in 2\mathbb{Z}_{\geq 1}$, $q \in \mathbb{Z}$.*

Proof. Assume that there exists a knot K' in $L(p,q)$ whose preimage is isotopic to $T_{3,2}$. By Theorem 3.1, $G(T_{3,2}) = \langle a, b \mid a^3 = b^2 \rangle$ is a C^p-group. We define a surjective homomorphism $\phi\colon G(T_{3,2}) \twoheadrightarrow \mathfrak{S}_3$ by $\phi(a) = (123)$, $\phi(b) = (12)$. One can check that $\operatorname{Ker}\phi$ is characteristic. Indeed, by [17], it suffices to show that $\phi(\theta(w)) = 1$ for any $w \in \operatorname{Ker}\phi$, where $\theta \in \operatorname{Aut}(G(T_{3,2}))$ is defined by $\theta(a) = a^{-1}$, $\theta(b) = b^{-1}$. Thus, $\mathfrak{S}_3$ is also a C^p-group by Lemma A.1. However, this contradicts Example 2.8. □

A.2. **The 6th symmetric group.** In this subsection, we prove that $\mathfrak{S}_6$ is not a C^p-group for even p (see Example 2.3). The following argument is based on [9, Theorem 3]. Let α denote the homomorphism $\mathfrak{A}_6 \hookrightarrow \mathfrak{S}_6 \rightarrowtail \operatorname{Aut}(\mathfrak{S}_6)$.

Lemma A.3 (see [9, p. 125])**.** $\mathrm{C}^p(\operatorname{Aut}(\mathfrak{S}_6)) = \operatorname{Im}\alpha$ *for $p \in 2\mathbb{Z}_{\geq 1}$.*

Proof. It follows from $[\mathfrak{S}_6, \mathfrak{S}_6] = \mathfrak{A}_6$ that $\mathrm{C}^p(\operatorname{Aut}(\mathfrak{S}_6)) \geq \operatorname{Im}\alpha$. One has to show that ϕ^p and $[\phi, \phi']$ belong to $\operatorname{Im}\alpha$ for any $\phi, \phi' \in \operatorname{Aut}(\mathfrak{S}_6)$. We prove only $\phi^p \in \operatorname{Im}\alpha$ for $\phi \notin \operatorname{Inn}(\mathfrak{S}_6)$. (The remainder is shown similarly.)

It is known that there exists $\psi \in \operatorname{Aut}(\mathfrak{S}_6) \setminus \operatorname{Inn}(\mathfrak{S}_6)$ whose order is 10. Since $\operatorname{Out}(\mathfrak{S}_6) \cong \mathbb{Z}_2$, we have $\psi^2 = \operatorname{Ad}_\tau$ for some $\tau \in \mathfrak{S}_6$. It follows from $\psi^{10} = \mathrm{id}_{\mathfrak{S}_6}$ that $\tau^5 = 1$, and thus $\tau \in \mathfrak{A}_6$. Hence, ϕ is written as $\phi = \psi \circ \operatorname{Ad}_\sigma$ for some $\sigma \in \mathfrak{S}_6$. Then we have

$$(\psi \circ \operatorname{Ad}_\sigma)^p = (\psi^2 \circ \operatorname{Ad}_{\psi^{-1}(\sigma)\sigma})^{p/2} = \operatorname{Ad}_{(\tau\psi^{-1}(\sigma)\sigma)^{p/2}}\,.$$

Since $\psi^{-1}(\sigma)\sigma \in \mathfrak{A}_6$, we conclude $\phi^p \in \operatorname{Im}\alpha$. □

Lemma A.4 (see [9, Theorem 3])**.** *For any C^p-group G, $\mathrm{C}^p(\operatorname{Aut}(G)) \geq \operatorname{Inn}(G)$, where $p \in \mathbb{Z}_{\geq 1}$.*

Proposition A.5. $\mathfrak{S}_6$ *is not a C^p-group.*

Proof. Assume $\mathfrak{S}_6$ is a C^p-group. By Lemma A.4, we have $\mathrm{C}^p(\mathrm{Aut}(G)) \geq \mathrm{Inn}(G)$, and thus

$$2 = [\mathrm{Aut}(\mathfrak{S}_6) : \mathrm{Inn}(\mathfrak{S}_6)] = [\mathrm{Aut}(\mathfrak{S}_6) : \mathrm{C}^p\mathrm{Aut}(\mathfrak{S}_6)][\mathrm{C}^p\mathrm{Aut}(\mathfrak{S}_6) : \mathrm{Inn}(\mathfrak{S}_6)].$$

On the other hand, we have $\mathrm{Aut}(\mathfrak{S}_6)/\mathrm{C}^p\mathrm{Aut}(\mathfrak{S}_6) \twoheadrightarrow \mathbb{Z}_2/\mathrm{C}^p\mathbb{Z}_2 = \mathbb{Z}_2$. Hence, $\mathrm{C}^p\mathrm{Aut}(\mathfrak{S}_6) = \mathrm{Inn}(\mathfrak{S}_6) \cong \mathfrak{S}_6$. This contradicts Lemma A.3. □

A.3. **Alternating groups.** We finally determine the group $\mathrm{C}^p(\mathfrak{A}_n)$.

Proposition A.6. *For $p \in \mathbb{Z}_{\geq 2}$ and $n \in \mathbb{Z}_{\geq 3}$,*

$$\mathrm{C}^p(\mathfrak{A}_n) = \begin{cases} 1 & \text{if } n = 3 \text{ and } 3 \mid p, \\ V_4 & \text{if } n = 4 \text{ and } 3 \mid p, \\ \mathfrak{A}_n & \text{otherwise,} \end{cases}$$

where V_4 is the Klein four-group.

Proof. The case of $n = 3$ is obvious since $\mathfrak{A}_3 = \mathbb{Z}_3$. In the case of $n \geq 5$, it is well known that $\mathfrak{A}_n^{\mathrm{ab}} = 0$, and thus $\mathrm{C}^p\mathfrak{A}_n = \mathfrak{A}_n$. Since $[\mathfrak{A}_4, \mathfrak{A}_4] = V_4$, $\mathrm{C}^p\mathfrak{A}_4$ is isomorphic to V_4 or $\mathfrak{A}_4$. Here, $\mathrm{C}^p\mathfrak{A}_4$ contains cyclic permutations of order three if and only if $3 \nmid p$. □

Appendix B. Taking C^p repeatedly

In this section, we discuss the derived p-series of a knot group. In [4], for a prime p, the derived p-series $\{G^{(n)}\}_{n\geq 0}$ of a group G is defined by

$$G^{(0)} := G, \quad G^{(n+1)} := \mathrm{C}^p(G^{(n)}).$$

We use this notation when p is not only a prime but also a positive integer. Here, determining the successive quotients $G^{(n)}/G^{(n+1)}$ and the intersection $G^{(\omega)} := \bigcap_{n\geq 0} G^{(n)}$ is a fundamental problem.

Proposition B.1. *Let G be a group with $G^{\mathrm{ab}} \cong \mathbb{Z}^{\oplus r}$. Then the abelian group $G^{(n)}/G^{(n+1)}$ surjects onto $\mathbb{Z}_p^{\oplus r}$.*

Proof. The abelianization of G induces the homomorphism $G^{(n)} \twoheadrightarrow (\mathbb{Z}^{\oplus r})^{(n)}$ for each n. Hence, we obtain $G^{(n)}/G^{(n+1)} \twoheadrightarrow (\mathbb{Z}^{\oplus r})^{(n)}/(\mathbb{Z}^{\oplus r})^{(n+1)} \cong \mathbb{Z}_p^{\oplus r}$. □

Remark B.2. In contrast, the stability of the (usual) derived series of $G(K)$ depends on a knot K ([5, Corollary 4.8], [15, Proposition 2.7]).

Proposition B.3. *Let p be a prime, G be a group with $H_1(G) \cong \mathbb{Z}^{\oplus r}$ and $H_2(G) = 0$. Then $G^{(\omega)}$ coincides with the commutator subgroup of G.*

Proof. Since the abelianization $\alpha\colon G \twoheadrightarrow \mathbb{Z}^{\oplus r}$ induces an isomorphism on $H_1(-;\mathbb{Z}_p)$ and a surjection on $H_2(-;\mathbb{Z}_p)$, by [4, Corollary 4.3], α induces an isomorphism $G/G^{(\omega)} \to \mathbb{Z}^{\oplus r}/(\mathbb{Z}^{\oplus r})^{(\omega)} = \mathbb{Z}^{\oplus r}$. Therefore, $G(K)^{(\omega)}$ coincides with $\operatorname{Ker}\alpha = [G(K), G(K)]$. □

Corollary B.4. *For p and G as in Proposition B.3, $G^{(n)}/G^{(n+1)}$ is isomorphic to $\mathbb{Z}_p^{\oplus r}$.*

Proof. We first have homomorphisms

$$G^{(n)}/\mathrm{C}^p(G^{(n)}) = G^{(n)}/G^{(n+1)} \hookrightarrow G/G^{(n+1)} \twoheadleftarrow G/G^{(\omega)} \cong \mathbb{Z}^{\oplus r}.$$

It follows that $G^{(n)}/G^{(n+1)} \cong \mathbb{Z}_p^{\oplus r'}$ for some $r' \leq r$. By Proposition B.1, we conclude $r' = r$. □

Remark B.5. Even if p is not prime, by considering a prime factor of p, we have $G^{(\omega)} \leq [G, G]$. On the other hand, if $6 \mid p$, then the equality fails for $G = G(3_1)$, where 3_1 is the trefoil. Indeed, since $\mathfrak{S}_3^{(2)} = \mathfrak{A}_3^{(1)} = 1$, the homomorphism $\phi\colon G(3_1) \twoheadrightarrow \mathfrak{S}_3$ defined in the proof of Theorem A.2 induces $G(3_1)/G(3_1)^{(\omega)} \twoheadrightarrow \mathfrak{S}_3$. Since $\mathfrak{S}_3$ is not abelian, $G(3_1)^{(\omega)}$ is strictly smaller than the commutator subgroup of $G(3_1)$.

The same argument shows that if $G(K)$ surjects onto $G(3_1)$, then the equality fails as well. Such knots are found, for example, in [10, Theorem 1.2].

References

[1] M. Boileau and E. Flapan. Uniqueness of free actions on S^3 respecting a knot. *Canad. J. Math.*, 39(4):969–982, 1987.

[2] G. Burde and H. Zieschang. Eine Kennzeichnung der Torusknoten. *Math. Ann.*, 167:169–176, 1966.

[3] N. Chbili. A new criterion for knots with free periods. *Ann. Fac. Sci. Toulouse Math. (6)*, 12(4):465–477, 2003.

[4] T. Cochran and S. Harvey. Homology and derived p-series of groups. *J. Lond. Math. Soc. (2)*, 78(3):677–692, 2008.

[5] T. D. Cochran. Noncommutative knot theory. *Algebr. Geom. Topol.*, 4:347–398, 2004.

[6] P. E. Conner and F. Raymond. Manifolds with few periodic homeomorphisms. In *Proceedings of the Second Conference on Compact Transformation Groups (Univ. Massachusetts, Amherst, Mass., 1971), Part II*, pages 1–75. Lecture Notes in Math., Vol. 299. Springer, Berlin, 1972.

[7] R. Hartley. Knots with free period. *Canad. J. Math.*, 33(1):91–102, 1981.

[8] R. Hartley and K. Murasugi. Homology invariants. *Canad. J. Math.*, 30(3):655–670, 1978.

[9] M. Haugh and D. MacHale. The subgroup generated by the squares. *Proc. Roy. Irish Acad. Sect. A*, 97(2):123–129, 1997.

[10] K. Horie, T. Kitano, M. Matsumoto, and M. Suzuki. A partial order on the set of prime knots with up to 11 crossings. *J. Knot Theory Ramifications*, 20(2):275–303, 2011.

[11] A. Kawauchi, (ed.). *A survey of knot theory*. Birkhäuser Verlag, Basel, 1996. Translated and revised from the 1990 Japanese original by the author.

[12] K. Kodama and M. Sakuma. Symmetry groups of prime knots up to 10 crossings. In *Knots 90 (Osaka, 1990)*, pages 323–340. de Gruyter, Berlin, 1992.

[13] E. Manfredi. Lift in the 3-sphere of knots and links in lens spaces. *J. Knot Theory Ramifications*, 23(5):1450022, 21, 2014.

[14] D. J. S. Robinson. *A course in the theory of groups*, volume 80 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1996.

[15] S. K. Roushon. Topology of 3-manifolds and a class of groups, 2002, arXiv:math/0209121.
[16] M. Sakuma. Uniqueness of symmetries of knots. *Math. Z.*, 192(2):225–242, 1986.
[17] O. Schreier. Über die gruppen $A^aB^b = 1$. *Abh. Math. Sem. Univ. Hamburg*, 3(1):167–169, 1924.
[18] J. Stallings. Homology and central series of groups. *J. Algebra*, 2:170–181, 1965.
[19] H. S. Sun. On groups generated by the squares. *Fibonacci Quart.*, 17(3):241–246, 1979.

Graduate School of Mathematical Sciences, the University of Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo, 153-8914, Japan
E-mail address: nozaki@ms.u-tokyo.ac.jp